


ΕΥΡΩΠΑΪΚΟ ΚΕΝΤΡΟ ΓΙΑ ΤΗ ΠΡΟΦΥΛΑΞΗ ΚΑΙ ΠΡΟΒΛΕΨΗ ΣΕΙΣΜΩΝ
ΑΘΗΝΕΣ, ΕΛΛΑΔΑ

ΕΛΛΗΝΙΚΗ ΕΤΕΡΩΠΕΙΑ
Ι. ΦΟΥΝΤΟΥΛΗΣ

ΕΥΡΩΠΑΪΚΟ ΚΕΝΤΡΟ ΓΙΑ ΤΗ ΠΡΟΦΥΛΑΞΗ ΚΑΙ ΠΡΟΒΛΕΨΗ ΣΕΙΣΜΩΝ
ΑΘΗΝΕΣ, ΕΛΛΑΔΑ

ΕΤΕΡΩΠΕΙΑ ΚΑΙ ΠΡΟΤΕΚΤΟΡΙΑ ΟΡΓΑΝΙΣΜΟΣ
ΑΘΗΝΕΣ, ΕΛΛΑΔΑ


SEMINAR ON ACTIVE FAULTS



September
14-17, 1995
ATHENS-PELOPONNESE

Excursion leader:
Prof. I. MARIOLAKOS

FIELDGUIDE



Pre-Congress Excursion A1 of the
XV Congress of the Carpatho-Balkan Geological Association

LEKKAS, E., PAPANIKOLAOU, D., FOUNTOULIS, I., (1995). - The Pyrgos earthquake.- The geological and geotechnical conditions of Pyrgos area (W. Peloponnese, Greece). *In Field-guide for the Pre-Congress Excursion of the XV Congress of the Carpatho-Balkan Geological Association, Athens, Sept. 1995*, pp. 42-46.

SEMINAR on ACTIVE FAULTS

Athens, Peloponnese, September 13-17, 1995

ORGANIZING COMMITTEE

Scientific Coordinator of the Seminar	:	Prof. I. MARIOLAKOS (Univ. Athens)
President of E.C.P.F.E.	:	Prof. D. PAPANIKOLAOU (Univ. Athens)
Secretary	:	Dr. I. FOUNTOULIS (Univ. Athens)
Cashier	:	Mrs H. MACRIS (EPPO)
Member	:	Mrs G. PAGONI (EPPO)
»	:	Drs A. ADAMOPOULOU (Univ. Athens)
»	:	Drs P. KARVELIS (Univ. Athens)
»	:	Drs P. VASSILOPOULOU (Univ. Athens)
*Excursion Leader	:	Prof. Ilias MARIOLAKOS

The Organizing Committee would like to thank the following Post-graduate students for their assistance in the preparation of the fieldguide for the Fieldtrip:

- Kranis H.
- Theoharis D.
- Zeri S.
- Badekas I.
- Tzavaras M.

THE PYRGOS EARTHQUAKE - THE GEOLOGICAL AND GEOTECHNICAL CONDITIONS OF THE PYRGOS AREA (W. PELOPONNESE, GREECE)

by

E. LEKKAS*, D. PAPANIKOLAOU*, I. FOUNTOULIS*

INTRODUCTION

The tectonic graben of Pyrgos, which extends over a large area, is characterized by complex lithostratigraphical structure and by neotectonic deformation which has been particularly intense during Holocene (HATZFELD et al., 1990, LEKKAS et al., 1992).

However, the prevailing conditions at the Pyrgos pleistoseismal area are relatively simple. Neotectonic structures are covered by recent geological formations (Alluvial, Marshy deposits) which occur in the flat low land.

On March 26 1993 the major area of Pirgos was affected by an earthquake of magnitude Ms:5.2 (Fig. 1).

The main shock as well as the aftershock caused many damages at Pirgos and the surraending area.

The Pirgos earthquake, except the destructions to the buildings caused several other geodynamic phenomena such as liquefaction phenomena followed by and water's sharing off, landslides, fractures, soil fractures etc.

Seismic fractures

Seismic fractures were observed at the following two areas.

A. Pirgos: Several seismic fractures, were observed at the northeastern part of the city in a general direction E-W. They have en chelon arrangement. Their length varies from some meters up to 10 m, the opening (width) is 2-5 cm. Any vertical throw is not visible. On the other hand there is well defined right lateral component of the movement.

The seismic fractures cut the asphalt of the road as well as some particles of the coarse-grained clastic rock in the asphalt. It must be mentioned that in the area which seismic fractures were observed, the damages were more (or) dense even in new constructions.

B. Lastetra village: Seismic fractures were observed at the southern part of the village. Their strike is E-W and have en chelon arrangement. Their length's varies from some centimeters up to some meters. They cut the asphalt of the road as well as the soils of the cultivated area. In some the building damages were connected by these seismic fractures.

GEOLOGICAL GEOTECHNICAL CONDITIONS

The approach to the geological and the accompanying geotechnical conditions of the city of Pyrgos has been carried out through the use of existing data of previous extensive researches and mainly the

* Univ. of Athen, Dept. of Geology
Dynamic Tectonic Applied Geology Div.
Panepistimiopolis Zografou 157 84
ATHENS GREECE.




Fig. 1. Block-diagram depicting the geological and neotectonic structure of Pyrgos area.




Fig. 2. Geological and tectonic map of Pyrgos area.

Neotectonic Map of Pyrgos area. (LEKKAS et al., 1992). This research has formed the source of data and information concerning the nature of the formations that occur in the graben, as well as the active macrostructure, a part of which is at Pyrgos city. For some of these, the determination of their geotechnical parameters was made possible (Fig. 3).

Geological mapping (Fig. 2) in the Pyrgos and the surrounding area, where identified the following formations described in reverse age order.

Recent Marshy Deposits. They develop surficially over the other formations at the flat plane part of the area and overlie mainly the Alluvial formations, although the contact between them is not well defined very clear. The most important outcrops overlay are north-west of Lapato and east of Aghios Athanasios quarters. They are alternations of brown-greyish-brown clays, grey, blue-grey clay silts, clay and silty sands containing abundant organic remnants. Total thickness is up to 5 meters, approximately. The Standard Penetration Test (S.P.T.) showed that the number of impacts is no more than 10.

Alluvial Formations. They occur at the flat area of the town of Pyrgos and overlay unconformably the older formations. They comprise of brown to grey soft clays with irregular (both in vertical and lateral sense) intercalation of brown silt and grey-brown sand. They contain numerous floral remnants as well as coarser material (gravel and pebbles). Their thickness does not exceed 12 m and the SPT test showed that the number of impacts is no more than 15.

Erymanthos Formation. It outcrops over a limited area, mainly at Kokkinohoma, Kouvelos and at the Aghioi Pantēs cemetery. It is a fossil outcrop of the Pleistocene Erymanthos formation, which comprises mainly polygenetic conglomerates of terrestrial origins connected with a red-siliceous fine unconsolidated formation. It corresponds to a huge paleo-talus cone with frequent lateral diversification and covers a large part of the graben of Pyrgos. At the research area in the city of Pyrgos it is represented by red to brown red clays and yellow brown sandy clays, loose horizontal sandy conglomerates and micro-conglomerates. It overlies unconformably on the Vounargos formations and its thickness varies from 2 to 8 metres. The SPT test showed that the number of impacts for a 30 cm penetration is no more than 15.




Fig. 3. Schematic lithostratigraphic column of all formations and results of the Standard Penetration Test.

Vounargos Formation. It is perhaps the most characteristic formation that outcrops in the graben of Pyrgos. Its age is Plio-Pleistocene, its thickness is up to 600 meters and comprises continuous intercalation of clays, silts, sandstones, sands and marls with constant diversification both in vertical and lateral sense. At the study area it occurs in the urban region and more specifically at the quarters of Aghia Kyriaki, Agios Nikolaos, and Aghios Athanasios. There can be distinguished two members of the formation which have both different lithological composition and geotechnical properties.

The upper member (Aghios Athanasios) develops at the quarters of Aghios Athanasios, Keramidhaki, Aghios Spyridon and partially at Lapato and practically covers all the lower parts of the uplands of Pyrgos. It comprises mainly yellow-brown cross-bedded sands and yellow silty sands with certain intercalation of yellow-brown sandy clays. At the SPT test the number of impacts for the clay sands and the silty sands is 15-30, while for the sandy clays it is 10-20. Its thickness exceeds 30 metres.

The lower member (Aghios Nikolaos) occurs at the quarters of Aghios Nikolaos and Aghia Kyriaki and covers most of the uplands of the city of Pyrgos. It comprises fossil bearing grey-blue-greyish marls which alternate with silty, sandy and clay marls, while locally there are thin intercalation of sand, sandy-silt, lignite horizons and yellow-brown sandstones. In contrast with the upper member, the SPT test showed that the number of impacts for 30 cm penetration exceeds 30, and sometimes they are more than 50. Its thickness is more than 80 m.

The formations that occur at the area of Pyrgos have undergone neotectonic deformation and are crossed by a number of faults of East-West mean direction. These faults are part of the Katakolo-Pyrgos fault zone, according to the existing literature. This fault zone was responsible for the earthquakes of March 26, 1993.

More specifically, the elongated outcrop of the lower member of Agios Nikolaos is abruptly terminated by the F1 (E-W bearing) fault, south of which the upper (Aghios Athanasios) member of the Vounargos formations occurs. It is a normal fault, accompanied by a morphological discontinuity (to the south of the city main square), its throw is at least 50 metres (estimated from morphotectonic features) and eastwards it branches into two faults. Existence of this fault is also confirmed by small polished surfaces.

To the north there is an identical picture with the occurrence of the F2 fault (av. direction NE -SW), which also branches into two minor faults to the east. The F2 fault brings together the outcrops of Aghios Athanasios and Aghios Nikolaos members, while it also crosses some outcrops of Erymanthos formation. Its throw is smaller, (20-30 metres) and all along it, we recognised seismic fractures caused by the shock of March 26, 1993.

One may distinguish, at the area of Pyrgos, an elongated horst of general East-West direction, which is pronounced by the ridge of the same direction. It consists of the lower (Aghios Nikolaos) member of

the Vounargos formation, as well as of a small part of Erymanthos formation. To the north and south of the horst there are the outcrops of the younger formations, which are represented by Aghios Athanasios member (it belongs to the Vounargos formation) and Holocene deposits (Alluvial and Marshy deposits).

DISCUSSION - CONCLUSIONS

Based on the intensity distribution map and the geological map of the urban complex of Pyrgos, we may propose the following basic correlations and results:

High intensities are observed alongside and on either side of F1 and F2 that cross the geological formations. The VIII isoseismals of the E.M.S. - 1992 present an impressive elongated development that also holds for the lower intensity coseismals. As already mentioned, these faults belong to the Katakolo - Pyrgos - Epitalio fault zone¹ parts of which were reactivated during the March 26, 1993 earthquakes. Along this fault zone, and more precisely along the F2 fault, seismic fractures were observed and those were the locations where the damages were particularly intense. A similar diversification of the destruction was observed along another fault zone at Kastro village near Pyrgos area during the shock of October 16, 1985.

Particularly high intensities were observed at the areas of Kouvelos and Kokkinohoma, where the fossil outcrops of Erymanthos formation overlie unconformably the Vounargos formation. The low geotechnical properties of these outcrops and their small thickness seem to have been the crucial factors for the magnification of intensities, having created a disadvantageous geodynamic regime.

Some formations or parts of them have caused local isolated "islets" of high intensity coseismals north-west of Lapato and east of Aghios Athanasios, where Marshy deposits with poor geotechnical properties, occur. In addition are also observed high intensities, west of the Aghioi Pantos cemetery. They coincide with the presence of loose sands that belong to Aghios Athanasios member of Vounargos formation. At this very location, even antiseismic designed buildings (e.g. schools) underwent severe damages.

REFERENCES

ΑΡΜΙΣΟ

- VARMISO, R., LYON-CAEN, H., PAPANASTASIOU D. (1992). East-west extension and Holocene normal-fault scarps in the Hellenic arc *Geology*, v. 20 p. 491-494.
- ΑΡΜΙΣΟ, R., LYON-CAEN, H., PAPANASTASSIOU D. (1991). A possible normal-fault rupture for the 464 B.C. Sparta earthquake. *Nature*, v. 351 p. 137-139.
- BOUKOUVALAS G. - SABATAKAKIS, N. 1987. Soil conditions and building damage in Kalamata from the 1986 earthquakes: *Bull. of the publ. works res. cent.*, Vol. 4, Oct.-Dec. 1987, pag. 267-275 (In Greek).
- CHRISTODOULOU, G., (1969). - Γεωλογικός χάρτης της Ελλάδας κλ. 1:50.000 Φύλλο "Βαρθολομιά", ΙΓΜΕ, Αθήνα.
- DEWEY, J.F. & SENGOR, C.A.M. (1979). Aegean and surrounding regions: complex multiplate and continuous tectonics in a convergent zone. - *Bull. geol. Soc. Amer.* 90: 84-92: Boulder.
- DRAKOPOULOS, J. & DELIBASIS N., 1982. The focal mechanism of earthquakes in the major area of Greece for the period 1947-1981. *Seismol. Lab. Univ. Athens. Publ.* 2, 27p.
- FOUNTOULIS, I. (1994) - Neotectonic evolution of Central - Western Peloponnesus Ph. D. Thesis Univ. of Athens, Dept of Geology, 254. p.
- FREYBERG, B.Y. (1973) - *Geologie des Isthmus von Korinth*. *Frlanger geol. Abh. Heft* 95; 183 Seiten.
- GALANOPOULOS, A. 1972. Plate tectonics in the area of Greece, as reflected in the deep focus seismicity. *Bull. Geol. Soc. of Greece*, IX, 2, 266-285.
- GEORGADAS, D. & LAGIOS, E. (1982): Remeasurement of the National Gravity Base Network in Peloponnesus. - *Ann. geol. Pays hellen.*, 31: 136-145; Athenes.
- HATZFELD, D., PEDOTTI, G., HATZIDIMITRIOU, P., MAKROPOULOS, K., (1990). The strain pattern in the western Hellenic arc from a microearthquake survey. *Geophys. J.*, vol. 101, 1, p. 181-202.
- KELLETAT, D. & SCHRODER, B. (1975): Vertical displacement of Quaternary shoreline in the Peloponnese (Greece): - *S.I.E.S.M., Rap. comm. int. Mec. Medit.*, 23/4a: 199-200.
- KELLETAT, D., KOWALCZYK, F., SCHRODER, B., WINTER, K.P. (1978): Neotectonics in the Peloponnesian Coastal Regions. In: CLOSS, H. et al. (Eds): *Alps, Apennines, Hellenides*: 512-518.
- KOWALCZYK, G. & WINTER, K. (1979). - Outline of the Cenozoic history of the Kylini peninsula, W. Peloponnese. In: Symeonides, N. Papanikolaou D. and Dermitzakis, M. *Field guide to the Neogene of Megara - Peloponnesus - Zakynthos* Department of Geology and Palaeontology, S.A. No 34.
- LAJ, G., JAMET, M., SOREL, D., & VALENTE, S.R., 1982. First pale tomagnetic results from Miocene series of the Hellenic sedimentary arc. *Tectonophysics*, 86, 45-6?
- LEKKAS, E., & DANAMOS, G., 1989. Impact of the geological conditions on the distribution of the damages at Kastro village (Kylini peninsula) caused by the earthquake of Oct. 16, 1988. *Bull. of the Geol. Soc. of Greece*, XXIV, (in press) (In Greek).
- LEKKAS, E., PAPANIKOLAOU, D., & FOUNTOULIS, J., 1992. Neotectonic map of Greece, Sheet "Pyrgos" - "Tropaia" (scale 1:100.000). Project, University of Athens, Dynamic, Tectonic, Applied Geology Div., 120p (in Greek).
- LEKKAS, E., PAPANIKOLAOU, D., FOUNTOULIS, J., & DANAMOS, G., 1994. Tectonic analysis of fault at the earthquake-stricken area of Pyrgos. Project, University of Athens, Dynamic, Tectonic, Applied Geology Div., 90p. (in Greek).
- LE PICHON, X. & ANGELIER, J. 1979. The Hellenic Arc and Trench system: a key to the neotectonic evolution of the Eastern Mediterranean area. *Tectonophysics* 60, 1-42.
- LE PICHON et al, 1979. From subduction to transform motion. A seabeam survey of the Hellenic trench system. *Earth & Plan. Sc. Let.*, 44, 441-450.
- LE PICHON, X. & ANGELIER, J. 1981. The Aegean Sea. *Phil. Trans. R. Soc. London*, A 300, 357-372.
- MARIOLAKOS, I. (1974): Comparative geomorphological observations between the drainage patterns of Erymanthos and Ladon (Peloponnesus, Greece). - *Prakt. Acad. Athens*, 49: 238-250.

- MARIOLAKOS, H., 1976. Σκέψεις και απόψεις επί ορισμένων προβλημάτων της Γεωλογίας και Τεκτονικής της Πελοποννήσου. *Ann. Geol. Pays Hellen.*, 27, 215-313.
- MARIOLAKOS, I., LEKKAS, S. & PAPANIKOLAOU, D., (1976): Quantitative geomorphological analysis of drainage patterns in the Vth order basins of Alfios River (Peloponnesus, Greece. - *Ann. geol. Pays hellen.*, 30, 2: 441-454, Athenes.
- MARIOLAKOS, I. & PAPANIKOLAOU, D. (1981a): The influence of the map scale on the results of the quantitative geomorphological analysis exemplified by Alfios River (Peloponnesus, Greece - *Ann. geol. Pays hellen.*, 30, 2: 441-454, Athenes.
- MARIOLAKOS, I. & PAPANIKOLAOU, D. (1981b): The Neogene Basins of the Aegean Arc from the paleogeographic and the geodynamic point of view. - *Proc. int. Symp. Hellenic Arc and Trench, Athens 1981: 383-399.*
- MARIOLAKOS, I., SYMEONIDIS, N., LEKKAS, S., KAROTSIERIS, Z. & SIDERIS, CH. (1981): The deformation of the area around the eastern Corinthian Gulf affected by the earthquakes of February to March 1981. - *Proc. int. Symp., Hellenic Arc and Trench, Athens 1981: 400-420.*
- MARIOLAKOS, I., PAPANIKOLAOU, D. (1985): Deformation pattern and relation between deformation and seismicity in the Hellenic arc. - *Bull. Geol. Soc. Greece, XIX, p. 59-76 (In Greek).*
- MARIOLAKOS, I., PAPANIKOLAOU, D. & LAGIOS, E. (1985). A neotectonic geodynamic model of Peloponnesus based on morphotectonics, repeated gravity measurements and seismicity. *Geol. Jb.*, B-50, 3-17.
- MARIOLAKOS, I., SABOT, E., LOGOS, E., LOZIOS, S., MERTZANIS, A., FOUNTOULIS, J. 1987a. The Geographical distribution of the rockfalls caused by the earthquakes of Kalamata. *Proceedings 1st Geogr. Congr., Athens 1987, v.B, p. 119-133 (In Greek).*
- MARIOLAKOS, I., SABOT, V., LOZIOS, S., LOGOS, E., FOUNTOULIS, J. 1987 Morphotectonic observations at the graben of the Dimiova-Perivolakia area. *Proceedings 1st Cong. of the Geogr. Soc. of Greece, Athens 1987, v.B, p. 119-133 (In Greek).*
- MARIOLAKOS, I., LOGOS, E., LOZIOS, S., FOUNTOULIS, J. 1988 Neotectonic deformation of the Zimbeli Fault surface (east of Kalamata town) town. *Proceedings 4th Congr. of the Geol. Soc. of Greece., Athens May 1988, v. XXIII/3, p. 241-258 (In Greek).*
- MARIOLAKOS, I., FOUNTOULIS, J., LOGOS, E., LOZIOS, S. 1989. Surface faulting caused by the Kalamata - Greece earthquakes (13.9.86). *Tectonophysics, Vol. 163, p. 197-203.*
- MARIOLAKOS I., LEKKAS, E., DANAMOS G., LOGOS, E., FOUNTOULIS I., ADAMOPOULOU, E. (1990) - Neotectonic evolution of the Kyllini Peninsula (NW Peloponnesus). *Proceedings of the 5th Congress, Thessaloniki, May 1990, Bul. Geol., Soc. Greece vol. XXV/3, p. 163-176, 1991 (In Greek).*
- MARIOLAKOS I., DANAMOS, G., FOUNTOULIS, I., LEKKAS E., LOGOS, E., (1991). Soil fractures and sand water's shaking off observed during the earthquake of October 16th, 1988 at the region of Vartholomio (W. Peloponnesus, Greece). In *Proceedings of the European School of Climatology and Natural Hazards Course* (Editors: M.E. Almeida - Teixeira, R. Fartechi, R. Oliveira, A. Gomes Coelho), *Com. Europ. Commun. EUR 12918 EN, p. 257-265, Brussels.*
- MARIOLAKOS, I., FOUNTOULIS, I., LOGOS, E., LOZIOS, S., (1991) - Methods to study the forssional neotectonic deformation: the case of Kalamata area (SW Peloponnesus, Greece). *Proceedings of IGCP project 250 Edit chen Qingxuan, Institute of Geomechanics, CAGS, v.3, p. 15-21, printed by Seismological Press.*
- MARIOLAKOS, I., SCHNEIDER, H., FOUNTOULIS, I., VOULOUMANOS, N., 1992 - Paleogeography. Sedimentation and neotectonic implications at the Kambos depression and Kitries day area (Messinia Peloponnesus (Greece). *Proceedings of the 6th Congress of the Geological Society of Greece. Athens, May 1992 (in print).*
- MARIOLAKOS, I., BADEKAS, I., FOUNTOULIS, I., THEOCHARIS, D., (1994). Reconstruction of the Eacly Pleistocene paleoshore and paleorelief os SW Peloponnesus area. *Proceedings of the 7th Congress of Geol. Society of Greece. May 1994 (In print).*
- MCKENZIE, D.P. 1970. Plate tectonics in the Mediterranean Region. *Geoph. J.R. astr. soc.*, 30, 109-185.

- McKENZIE, D.P. 1978. Active tectonics of the Alpine-Himalayan belt the Aegean sea and surrounding regions. *Geoph. J.R. astr. soc.*, 55 (1), 217-254.
- MERCIER, J.L. 1979. Signification neotectonique de l' Arc Egeen. Une revue des idees. *Rev. Geol. Dyn. Geogr. Phys.* 21, 1 5-15.
- PAPANIKOLAOU, D. 1984. Introduction to the Geology of Greece. IGCP Proj. No 5, Field meeting in Greece. Sept. 17-23 1984, Field Guide, 3-35.
- PAPANIKOLAOU, D. 1985. The three metamorphic belts of the Hellenides: a review and a Kinematic interpretation. The Geological evolution of the Eastern Mediteranean. Publ. of the Geol. Soc., No 17, Blackwell Scientific Publ., Oxford 848 pp.
- PAPANIKOLAOU, D. 1986, Geology of Greece. University of Athens, 240 p. (in Greek).
- PAPANIKOLAOU, D. & DERMITZAKIS, M. (1981): The Aegean Arc during Burdigalian and Messinian: a comparison - *Riv. ital. Paleont.* 87, 1: 83-92, Milano.
- PAPAZACHOS, B.C., KIRATZI, A.A., HATZIDIMITRIOU, P.M. & ROCCA, A.C. 1984. Seismic faults in the Aegean area. *Tectonophysics*, 106, 71-85.
- PAQUIN, C. FROIDEVEAUX, C., BLOYET, J., RICARD, Y. & ANGELIDIS, C. 1982. Tectonic stresses on the mainland of Greece: in-situ measurements by overcoring. *Tectonophysics*, 86.
- POULIMENOS, G., ZELILIDIS A., KONTOPOULOS, N. and DOUTSOS Th. - 1993 - Geometry of trapezoidal fan deltas and their relationship to extensional faulting along the southwestern active margins of the Corinth rift, Greece. *Basin Research* 5, p. 172-192.
- PHILLIPSON, A. 1930. Beitrage zur Morphologie Griechelands. *Geogr. Ab h.* 3, 1-96.
- RITSEMA, A.R. 1974. The earthquake mechanisms of the Balkom region. *Roy. Netherl. Meterol. Inst., De Bilt, Scient. Rep.*, 74-4, 36p.
- WHITCOMB, J.H. (1976): New Vertical Geodesy. - *J. geophys. Res.*, 81: 4937-4944.