

INTERNATIONAL ASSOCIATION OF ENGINEERING GEOLOGY



INTERNATIONAL SYMPOSIUM ENGINEERING GEOLOGY and the ENVIRONMENT

Engineering Geology Committee of the Geological Society of Greece
(Greek National Group of IAEG)



CORINTH CANAL, CORINTHIAN GULF, DELPHI AND CENTRAL GREECE

Editors: I. Mariolakos and I. Fountoulis, Geology and Tectonics
P. Marinos, G. Tsiambaos and M. Kavvadas, Engineering Geology
and Geotechnical Engineering, Hydrogeology

BOAT AND FIELD TRIP, WEDNESDAY, JUNE 25, 1997

ATHENS, GREECE, JUNE 23-27, 1997

MARIOLAKOS, I., FOUNTOULIS, I., KRANIS, H., (1997). - *Introduction to the Geology of Sterea Hellas, Hellenic Territory: Current geodynamic regime, Neotectonics of Corinthiakos, gulf, Delphi, Minyan Ancient Flood Prevention works.* In boat and field trip guide of the 7th International Symposium Engineering Geology and the Environment IAEG (Editors I. Mariolakos and I. Fountoulis, Geology and Tectonics ,P. Marinos,G.Tsiambaos and M. Kavvadas Engineering geology and Geotechnical Engineering, Hydrogeology), 60 p.

International Association of Engineering Geology
 Association Internationale de Géologie de l'Ingenieur


International Symposium
 Symposium International

Engineering Geology
 and the Environment
 La Géologie de l'Ingenieur
 et l'Environment



Field Trip Map

By I. Mariolakos, I. Fountoulis & H. Kranis



MAIN ISSUES OF THE FIELD TRIP

- Corinth Canal
- Corinthian Gulf
- Karstic masses of Central Greece
- Delphi
- The drained Kopaida lake

Legend of field trip map showing route and generalized geological features

1. Neogene-Quaternary deposits
2. Alpine clastic formations
3. Ophiolites
4. Alpine carbonates
5. Metamorphic rocks
6. Plio-Quaternary volcanic rocks
7. Major fault zones, visible during route
8. *Soussaki* volcano
9. Field trip route
10. Stop

CONTENTS

I. GEOLOGY AND TECTONICS

1. Introduction to the Geology of Sterea Hellas
2. Hellenic Territory: Current Geodynamic Regime
3. Saronikos Gulf
4. Neotectonic evolution of the Isthmus of Corinthos
5. Ancient Corinthos and Diolkos
6. Neotectonics of Corinthiakos Gulf
7. Delphi
8. The drained Kopaida Lake: Minyan Ancient Flood Prevention Works (c. 1600 BC)

II. ENGINEERING GEOLOGY AND GEOTECHNICAL ENGINEERING OF THE ISTHMUS AND THE CANAL OF CORINTH

1. Geological description from an engineering point of view
2. Geotechnical properties of the Corinth Canal Marls
3. The stability of the slopes of the Canal
4. Geotechnical problems of the existing railway bridge
5. Foundation of the new highway bridge over the Canal
6. Foundation of the new railway bridge over the Canal

III. HYDROGEOLOGICAL ISSUES OF CENTRAL GREECE

1. Karstification and groundwater hydraulics of the interior of large calcareous massifs: The case of Giona mountain in Central Greece
2. Assesments of groundwater resources based on GIS in the Central Greece

IV. ENGINEERING GEOLOGY OF THE DELPHI SITE

1. Engineering geology and geotechnical engineering of the archaeological site of Delphi, Greece
2. Geotechnical modeling of the limestone cliffs above the Delphi archaeological stie

ANNEX

Compiling of water balance of the Boetian Kifissos

Studies and works of the 1st Ephoria for the preservation and protection of the monuments
at Delphi

I. GEOLOGY AND TECTONICS

I. MARIOLAKOS, I. FOUNTOULIS, H. KRANIS

University of Athens, Department of Dynamic Tectonic Applied Geology

1. Introduction to the Geology of Sterea Hellas
2. Hellenic Territory: Current Geodynamic Regime
3. Saronikos Gulf
4. Neotectonic evolution of the Isthmus of Corinthos
5. Ancient Corinthos and Diolkos
6. Neotectonics of Corinthiakos Gulf
7. Delphi
8. The drained Kopaida Lake: Minyan Ancient Flood Prevention Works (c. 1600 BC)

Introduction to the geology of Sterea Hellas

I. Mariolakos, I. Fountoulis & H. Kranis

*University of Athens, Department of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84, Athens, Greece*

Greece represents a very characteristic part of the Alpine System, known as the Hellenic Arc. It represents one of the major mountain chains of the Alpine System, resulted from the convergence between two continental plates: The Eurasian to the North and the African to the South.

The main feature of the Hellenic Arc is its morphotectonic direction, which is NNW-SSE along Continental Greece. From Kythera island to Crete island it takes gradually an E-W direction and from Dodecaniss up to Isparta and Antalya (Turkey trends to a NE-SW direction (**Fig. 1.1**)).

A large number of tectonic units, corresponding to individual nappes, exists within the Hellenic Arc, and the overall kinematics show a movement directed from the core of the arc in the Aegean Sea towards the periphery in the Ionian and Libyan Sea. Generally the more internal units override the more external ones.

- The main characteristics of the alpine units in the Hellenides, are the following:
 - The paleo-Alpine orogeny of Late Jurassic – Early Cretaceous
 - The Alpine orogeny which started in Late Eocene and culminated during Oligocene and Miocene times. Nevertheless plate tectonic movements with resulting orogenic processes are still in action, along the present Hellenic Arc and Trench System.
- The alpine tectonic units in Greece are separated in Metamorphic and non-Metamorphic.

The main tectonic units distinguished in Sterea Hellas and more precisely the area of the field-trip are the following (see geotectonic map):

Parnassos:	Neritic carbonate sequence (interrupted by 3-4 bauxite horizons) and flysch (Triassic – Eocene).
Western Thessaly Beotia:	Continuous sequence from Triassic to Eocene. It is the most internal unit of the Hellenides with continuous stratigraphic column.
Eastern Greece:	It is deformed twice during the paleo-Alpine and typical Alpine orogeny. Comprises a neritic Triassic – Jurassic sequence (Subpelagonian) overlain by the obducted ophiolites and the transgressive Upper Cretaceous limestones and the Eocene flysch on top.




Fig. 1.1 The Hellenic Arc and Trench System. KF: Kefalonia Fault, NAF: North Anatolian Fault, PT: Plini Trench; ST: Strabo Trench.

Hellenic Territory: Current Geodynamic Regime

I. Mariolakos, I. Fountoulis & H. Kranis

University of Athens, Division of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84 Athens

The present Hellenic Orogenic Arc is restricted at the southern part of the Hellenic territory, in contrast to all the previous ones, which extended throughout the whole length of the Hellenides.

During the Middle Miocene, a part of the Hellenic arc, still active today, was cut off from the Tethyan chain and since then it followed its own evolution. To the north, this part is bounded by the prolongation of the right-lateral Anatolian fault (**Fig. 1.1**). In the northern Aegean region, this fault coincides with the northern limit of the active part of the Hellenic arc, bounding an area termed “Aegean microplate” (McKenzie, 1970; 1972; 1978; Galanopoulos, 1972).

The present geometry of the Hellenic Arc has been developing since the Late Miocene. The back-arc basin and the volcanic arc are restricted in the Aegean plate region.

According to Le Pichon *et al.* (1981), the present geodynamic regime of the Hellenic arc is characterized by asymmetrical movement; along the Ionian trench, the subduction direction is NE-SW and the regime is pure compression, in accordance with the fault plane solutions while, in the Pliny and Strabo trenches, the direction of movement is composite, featuring a substantial sinistral NNE-SSW horizontal component (**Fig. 1.1**). In the back-arc area there are extensional structures also with significant horizontal component of movement.

Many geodynamic models have been proposed for the Hellenic arc and especially for Peloponnesos. These models accept that the latter under extensional stress field, accompanied by graben created by normal faulting in the back arc basin (Ritsema, 1974; McKenzie, 1978; Mercier, 1979; Le Pichon & Angelier; 1979; Dewey & Sengör 1979 and others).

Mariolakos & Papanikolaou (1981) suggested that marginal fault zones control the configuration of neogene basins (**Fig. 2.1a**). These fault zones create an asymmetry to basin morphology and sedimentation. According to them, the Hellenic arc is separated in three large parts (**Fig. 2.1b**). In part I the big fault zones have an E – W direction. In part II the direction is NW - SE and in part III the direction changes to NE - SW. This arrangement shows that only parts II and III have apparent dynamic relation to the Hellenic arc and trench system, while part I has its own peculiarity.

Data on the current deformation pattern of the Hellenic Arc have been provided by numerous researchers, as:

- *in situ* measurements of the stress field in shallow drillings (<10m),(Paquin *et al.*, 1982)
- paleomagnetic investigation of the Neogene and Quaternary sediments (Laj *et al.*, 1982)
- fault plane solutions (McKenzie 1972; 1978, Ritsema, 1974; Drakopoulos & Delibasis, 1982; Papazachos *et al.*, 1984.

Mariolakos & Papanikolaou (1987) combined the results of various geological, seismological and geophysical studies, and proposed a present (active) deformation model (**Fig. 2.2**).

Further data that assist in the interpretation of the current deformation regime of the Hellenic territory have been supplied by geodetic measurements (Billiris *et al.*, 1989) and the distribution of earthquake foci (**Fig. 2.3**).

More recent investigations from SW Peloponnesos (Messinia) (Mariolakos *et al.*, 1991) showed that the stress field responsible for the neotectonic-active deformation is that of rotational couple

(**Figs. 2.4; 2.5**) which caused not only brittle but also ductile deformation structures resulting from local transpression and transtension sectors.




Fig. 2.1 The neotectonic map of the main marginal fault zones of the post-alpine basins in the southern continental Greece (a) and the neotectonic fault pattern of the Hellenic arc (after MARIOLAKOS *et al.*, 1985).




Fig. 2.2 Analysis of the general stress field F on the faults of the three parts (I, II, III) of Fig. 2.1, into pure (σ) and shearing ($T\alpha$) components. In the upper left part of the picture the NW Peloponnessos region is analyzed (under magnification). The faults are developing with a substantial horizontal sinistral component and a dextral rotation of the part in between. The relatively non-seismic Cyclades region is dominated by NW-SE oriented faults. In these faults, the shear component ($T\alpha$) of the general stress field is minimum (after Mariolakos & Papanikolaou, 1987).




Fig. 2.3 Seismicity map for the period 1900-1995.




Fig. 2.4 Torsional deformation pattern between Dimiova-Perivolakia graben and Mt. Kalathion Horst (Messinia, SW Greece)(After Mariolakos *et al.*, 1991)




Fig. 2.5 Torsional deformation and related structures (After Mariolakos *et al.*, 1991).

Saronikos Gulf

I. Mariolakos, I. Fountoulis & H. Kranis

*University of Athens, Division of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84 Athens*

Saronikos Gulf is located between the peninsulas of Argolis to the west and Attiki to the east. It has a complicated morphology and is divided into a western and an eastern part by a very shallow N-S trending platform, part of which emerges as the islands of Methana, Angistri, Aegina, Salamina (**Fig. 3.1**) (Papanikolaou *et al.*, 1988). This N-S zone, separating the Western and Eastern Saronikos gulfs, comprises several outcrops of Plio-Quaternary age, representing the northern edge of the modern volcanic arc. The Western Saronikos Gulf includes two basins, the WNW-ESE trending Epidauros basin to the south (more than 400m deep) and the E-W trending Megara basin to the north, which is relatively shallow (less than 250m). The Megara basin is a tectonic graben bounded by E-W to ENE-WSW marginal faults with a throw of 400-500m. Between the Epidauros basin to the south and Megara basin to the north, an alternation of horsts and graben is observed, bounded by E-W to ENE-WSW trending faults with throws between 200 and 500 m..

The western part of Saronikos Gulf is evidently more active than its eastern one, judging from fault displacement, sediment distribution and the presence of recent volcanoes, which delineates the active western part from the relatively inactive eastern part.




Fig. 3.1 Neotectonic sketch map of the Saronikos Gulf (after Papanikolaou *et al.*, 1988).

Neotectonic Evolution of the Isthmus of Corinthos

I. Mariolakis, I. Fountoulis & H. Kranis

University of Athens, Division of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84 Athens

The major area of Corinthos consists of post-alpine formations. It represents a neotectonic graben, bounded by two marginal, E-W striking fault zones, namely the Gerania Mt. fault zone to the North and the of Onia Mt. f.z to the south (Field Trip Map; **Fig. 4.1a**).

The Isthmus of Corinthos is a narrow strip of land that connects the Peloponnessos to the Hellenic mainland and marks the easternmost limit of the Corinthiakos Gulf. It is located in the above mentioned graben, and consists of a succession of uplifted, well-exposed Pliocene marls unconformably overlain by several cycles of near-shore Pleistocene conglomerates.

Neogene and Quaternary sediments are cut by numerous, NE-SW and E-W striking normal faults. Some of these can be observed along the 6.3-km-long canal, cut at the end of the nineteenth century along the path of *Diolkos*, an ancient ramp used as a vessel transportation route connecting the Corinthiakos and the Saronikos Gulfs. Because these faults belong to two different sets and dip to the north and south, respectively (section A-A' in **Fig.4.2**), the Isthmus has been interpreted as a horst; i.e., as a simple, extensional feature (Philippson, 1890).

An alternative interpretation based on the mapping of the unconformity between the Pliocene and Pleistocene deposits (Eutyrrhenian) was given by Freyberg (1973). Detailed geologic mapping and borehole and geophysical data were used for the compilation of this surface, shown in **Fig. 4.3**. The deformation of this originally horizontal surface to its present-day complex shape is presumed to express the Quaternary deformation pattern in the area. Or, as Freyberg (1973) stated, "*The Isthmus...seems to be formed by anti-tilted fault-blocks. The northern system of the block-faulted area has its greatest height in the East and dips below sea-level towards the West ("Tieflage") while the system in the South acts reversely, and between them, a neutral zone, tilted to a lesser degree, exists*" (see **Fig.4.3**). Thus, a NW-SE section shows an apparent horst structure, whereas any section along a southwest-northeast axis would disclose a graben (**Fig.4.2**). The deformation of the Isthmus, the cover of which consists of clay and sand, does not favor brittle fracturing and is accommodated by numerous parallel, homothetic and antithetic normal faults, most of them active, which impart a pseudoplasticity to the crust in this area.

The area of the Isthmus is tectonically active. In the past 60 years, three earthquakes (1928, 1953, and 1981) have been associated with moderate to minor surface faulting (**Fig. 4.1a**; Mariolakis and Stiros, 1986).

Lithofaga molluscs and beachrock are exposed at a height of about 1.1 m above the present sea level at the ancient (2000-yr-old) harbor constructions of Leheon; they disclose emerging beaches that can be followed up to Corinthos. Freyberg (1973) reported rounded pottery sherds of undetermined age in this area (Neolithic to Roman) cemented in the conglomerates of former shorelines. The latter are disappearing east of Corinthos, while at Possidonia, beachrock covers some fourth century B.C. constructions of the *Diolkos*, and an ancient platform next to it is cemented to 0.75 m above sea level. Farther north, at Loutraki, the same, probably beachrock structure is at a height of 1.0 m. At the eastern exit of the canal, submerged ruins of the ancient harbor of Schoenous have been identified at the modern site of Kalamaki. There are also data showing that the westernmost portion of the Saronikos Gulf coast is submerging; the harbor of Kehreai (5 km south of Kalamaki) is the best known example.

The long-term (Quaternary) and short-term (late Holocene to present) data discussed above are summarized in **Fig. 4.4**.

According to Mariolakos & Stiros (1987), the Corinthiakos Gulf separates a bulging area (anticline?) to the south and a structural depression to the north, the Isthmus marking their easternmost limit and an area where the associated deformations are minimum and easily observable. The depression to the north may extend as far as the Ionian Islands, which are characterized by pre-Quaternary reverse faults and a Holocene or even older tilt antithetic to that observed in the northern part of the Isthmus (**Fig.4.5**).

This analysis presumes that back-arc compression is not confined to a narrow zone hardly covering the westernmost end of the Peloponnesos, where minor contraction features have been reported (Mercier *et al.*, 1979), but may extend as far as the isthmus. In the northern Peloponnesos, however, this compression should be mild, thus producing only a very long wavelength bulging. This is contradictory to previous concepts that east-west normal faults and focal mechanism solutions of shallow earthquakes signify that this area is under extension (Mercier *et al.*, 1979). However, these analyses fail to explain the differential vertical motions along the Corinthiakos Gulf, as well as the antithetic tilting of the Isthmus strata. Furthermore, the fact that the crust in the northern Peloponnesos is up to 2.5 times thicker than in the Aegean, to the east (Makris, 1978), may suggest that the former is an area of crustal shortening.

Tselentis and Makropoulos (1986) computed the deformation tensor in the Corinthiakos Gulf by using large ($M > 5.5$) earthquake fault-plane solutions, and showed that north-south extension is not the dominant mode of deformation in this area. Their data seem to corroborate some north-south as well as some east-west extension, but the corresponding values are very small relative to the dominating deformation (uplift and subsidence of the southern and northern coasts, respectively) and may be insignificant, as no standard deviations are computed. However, some north-south extension, as a consequence of the east-west back-arc compression, is likely to exist. East-west-striking normal faults may originate from this second-order extension, especially since older lines of weakness are reactivated (Sebrier, 1977). However, things must be more complicated, because normal faults and focal mechanisms contain some characteristic strike slip.

Hatzfeld *et al.* (1990) computed 16 focal mechanism solutions of shallow earthquakes that show N-S mostly normal faulting, while few of them are strike-slip. However almost none of the normal mechanisms corresponds to pure dip-slip, but most of them are oblique-slip.

The previous discussion suggests that the well-documented torsional deformation of the Isthmus is taken up by steep, parallel normal faults. Consequently, these normal faults, as well as the normal faults flanking the Corinthiakos Gulf, are not indicative of regional extension, as was previously believed.




Fig. 4.1 a: Location map. Main faults, seismic faulting (thick lines), leveling benchmarks common in 1969 and 1981 topographic, and locations of cross sections of Fig. 4.2 are shown (after Mariolakos and Stiros, 1987)

b: Ground-surface displacement for the period 1969-1981: Central part of Isthmus (between benchmarks 79 and 84) appears uplifted relative to near-coastal areas. Subsidence of benchmark 77 is consistent with the motion expected at the hanging wall of the faults that ruptured in 1953 and 1981, as well as with a general subsidence tendency along the Saronikos Gulf. During the intersurvey period, the only important earthquakes that affected the Isthmus were the 1981 events. These earthquakes were associated with normal faulting about 10 km north of the study area. Therefore, any motion observed is likely to be associated with local tectonics.

c: Elevation change of the Eutyrrhenian layer along the leveling route. The resemblance of Figure 1b and 1c, presumably showing the present-day and quaternary deformation of the Isthmus, respectively may reveal that the pattern of crustal deformation in this area has not changed since the Early Quaternary.




Fig. 4.2 Schematic cross sections of Isthmus (for locations see Figure 4.1). Neogene deposits are shaded, Quaternary deposits are blank. In sections B-B' and C-C' Quaternary cover is too thin to appear at the scale used. (Based on data of Freyberg, 1973).




Fig. 4.3 Perspective view of contour map of Eutyrrhenian layer height, after Freyberg (1973). Canal and leveling benchmark positions (bold numbers) are also shown (After Mariolakos and Stiros, 1987).




Fig. 4.4 Quasi-perspective block diagram of Isthmus deformation (not to scale). Arrows indicate torsional deformation. LE=Leheon; CO=Corinthos; PO=Possidonia; LOU=Loutraki; KA=Kalamaki; KE=Kenchreai; AT=Aghioi Theodori.




Fig. 4.5 Differential movements and morphological features in Corinthiakos Gulf combined with deformation of Isthmus. a: location map, b: Quaternary uplift rates from sampling locations (marked with 'x' in (a)); c: axial bathymetry of Gulf; d: Late Holocene shoreline displacement (points correspond to solid circles in (a)) (After Mariolakos and Stiros, 1987).

Ancient Corinthos and Diolkos

I. Mariolakos, I. Fountoulis & H. Kranis

*University of Athens, Division of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84 Athens*

A few kilometres west of the modern town, the Ancient town of Corinthos is founded partially on the Neogene formations and partially on the Quaternary deposits (Tyrrenian). The construction material is mainly pleistocene calcareous oolitic sandstone (known as *poros*) that outcrops near the city of Corinthos.

Diolkos was a ramp connecting Corinthiakos to Saronikos Gulfs, on which the ancient vessels were pulled from one side to the other in order to avoid the circumnavigation of Peloponnessos. (**Fig.5.1**). It should be noted that although the people who profited from the existing harbors of Lechaion on the Corinthiakos gulf side and Kechreae on the Saronikos Gulf side *Diolkos*. However, the passage of ships was becoming more and more difficult as commerce was being developed and warfare necessitated the building of larger vessels.

It was *Periandros* (600 B.C.), the Tyrant of Corinthos, who first conceived the idea to cut the Isthmus by constructing a canal. He soon, however, gave up when he faced the technical problems involved in such a construction. In addition he was afraid of the wrath of *Poseidon* to whom the Isthmus area was dedicated.

Three hundred years later, *Dimitrios the Besieger* planned to construct the canal, but was stopped by his engineers who, (as mentioned by Strabo), asserted that the sea level of the Corinthiakos Gulf was higher than that of the Saronikos Gulf and thus a canal opening could cause the flooding of the coasts of Egina.

Julius Caesar and *Calligula* considered again the question of opening a canal. About a century later when *Nero* came to Corinthos to attend the *Isthmia* festivities, he took the decision to undertake this project. Indeed, the opening started both from the Corinthiakos and the Saronikos Gulf sides. However, after his return to Rome and his death, all work ceased. *Herodes Atticus* who later attempted to continue the project, was ordered to stop. Several hundreds years later, another attempt by the Venetians was condemned to failure.

Following the liberation of Greece, a new study was made but the estimated cost was more than the newly formed State could afford. Finally, half a century later on March 29th, 1882 the official foundation ceremony took place. The completed canal was inaugurated on July 25th, 1893.

The expert advice of some of the authorities of the time was sought and *Ferdinand de Lesseps* was consulted. The final design of the cut and the execution of the entire work was undertaken and brought to completion by the Hungarian civil engineer *Bela Gerster*.

Isthmus Canal - Technical data
Total length 6.300 m of which 540 m comprise the outer port.
The slopes at their base are walled by stones for a length of 3.500 m.
Width at the bottom: 21 m.
Width at the surface of the sea: 24.6 m.
Depth (uniform): 8 m.
Max. height: 79 m.




Fig. 5.1 The Isthmus with Canal and Diolkos.

Neotectonics of Corinthiakos Gulf

I. Mariolakos, I. Fountoulis & H. Kranis

*University of Athens, Division of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84 Athens*

Corinthiakos Gulf is a narrow strip of sea separating Peloponnessos from the rest of continental Greece. According to most researchers it is an asymmetrical tectonic trough, the exact formation time of which is unknown; however it must be between Late Miocene and Early Pliocene. Neotectonic and more specifically the Quaternary deformation studies around the Gulf revealed that the faulting as well as the morphology of the coast is the result of rotation phenomena (Freyberg, 1951; 1973; Mariolakos, 1976; Mariolakos *et al.*, 1985). Furthermore, Kelettat & Schröder, (1975), and Mariolakos *et al.*, (1989; 1991; 1994), suggest that certain blocks at the southern Peloponnessos have been subjected to rotation.

The distribution of the pliocene-quaternary deposits along the coast areas of the Corinthiakos Gulf is almost unilateral. North of the Gulf they are nearly absent, with the exception of one occurrence near Itea, the *Agia Efthymia* lacustrine conglomerates, of about 40-50 m thickness. In contrast, the plio-quaternary sediments are abundant on the southern coast, where they reach a thickness of more than 1200 m. They occur from the sea level up to a height of about 1700 m. The facies of the younger deposits (mainly sandstones and conglomerates) is more or less continental with intercalations of marine beds (Dercourt, 1964; Ori, 1989).

Heezen *et al.*, (1965) conducted an investigation of the sea-floor relief in the Gulf and demonstrated that there are significant differences between the northern and southern parts of it (**Fig. 6.1**). These differences can be observed on the continental shelf as well as on the continental slope of both margins. The continental shelf of the Gulf along the southern central part is, in general, poorly developed, seldom exceeding 500 m. in width. In contrast, along the northern coast it is consistently wider, (maximum width ~9 km in the bays Antikyra and Krissaion). It is also quite steep on the southern side, as opposed to the northern side where it dips more gently; furthermore, the latter is more intensively dissected and is characterized by a micro-relief. This asymmetry is more pronounced in the central part of the Gulf, where the 'abyssal plain' of the Gulf lies, just opposite to the area where the pliocene-quaternary sediments are most pronouncedly uplifted (i.e. Mts. Ziria and Helmos, north Peloponnessos) and reach their maximum thickness.

Recent research (Papanikolaou *et al.*, 1997) has shown that the thickness of the sediments in the Gulf increases eastward, and such is the case for the throw of the faults that bound the Gulf.

The following principal features characterize the topography of the northern Peloponnessos (south of the Gulf).

- i) High altitudes (Mt. Ziria: 2376 m; Mt. Chelmos: 2341 m; Mt. Panahaikon: 1926 m). All these mountains rise at small distances from the present shoreline.
- ii) The step-like relief of the northern part of the Peloponnessos with long and narrow terraces, which is partly due to old abrasion that followed the deposition of recent marine sediments.
- iii) The valleys of the northern Peloponnessos are deeply incised and steep gradients are another characteristic of the thalwegs. The geomorphology of the northern margin is completely different and the area occupied by the mountains Parnassos and Giona consists of four erosion surfaces. (Maul, 1921; Philippson, 1930; Sweeting, 1967). The step-like arrangement of the relief, which is so characteristic of the Northern Peloponnessos topography, is absent here. Furthermore, with very few exceptions, the streams of the northern margin are not incised.

The active fault zones of the Gulf comprise *en échelon*, right-stepping E-W striking faults (Papanikolaou *et al.*, 1997), and the mean direction of the Gulf is the result of this fault pattern (**Fig. 6.2**). The southern margin of the Gulf has been uplifting during the Plio-Quaternary, a fact confirmed by the occurrence of the plio-quaternary conglomerate beds at elevations higher than 1600 m (**Fig. 6.3**). (Mt. Mavrovouni).

In the vicinity of the Isthmus of Corinthos, Philippson (1892) distinguished two fault systems: the Corinthian fault system at the northern side of the Isthmus and the Crommyonian at the southern side. Although both systems have the same strike the former are homothetic whereas the latter are antithetic. The normal character of the faults in the area of the Isthmus had led to the conclusion that the disruption was caused by tensional stresses. Thus Philippson (1892), who first studied the geologic section of the canal, regarded it as a horst. However, later investigations by Freyberg (1951, 1971) showed that the tectonic movements have a rotational character and therefore the normal faults, regardless of their dip direction relative to that of the faulted strata, have resulted from rotation around an almost horizontal axis.

Based on Freyberg's conclusions that faulting in the area of the Isthmus of Corinthos has resulted from rotation involving tectonic processes, Mariolakos (1977), accepted that the region between Sperchios River basin to the north and Mts. Ziria, Chelmos and Panachaikon south of the Gulf, can be divided into a number of neotectonic multi-blocks. These, on account of their specific tectonic behaviour, were called *tectonic dipoles*. Thus, the mountains of Lidorikion, Giona and Iti constitute one tectonic dipole, which was referred to as the Giona - Iti tectonic dipole, whereas the northern part of Peloponnessos makes up the Corinthian tectonic dipole (**Fig. 6.4**). As Mariolakos (1977) pointed out, the tectonic dipole concept is not identical to that of the tectonic block; in fact a tectonic dipole consists of more than one block and forms a mosaic structure, that is, a tectonic multi - block.

The characteristic property of the tectonic dipole is that it shows a differential movement, due either to different direction of movement or different velocity. The differential movement is more obvious near the extremities of the dipoles, the *poles*, one of which appears to rise, relative to the other, which appears to descend. Furthermore, it must be noted that despite the apparent independence of the block-components of the dipole, the final movement, if there is one, resulting from the combination of the partial movements throughout the dipole is such that the two poles attain different movements or velocities relative to each other.

The Giona-Iti tectonic dipole and the Corinthian tectonic dipole, are two independent tectonic units which, nevertheless, behave in a similar manner in as much as there is an upward motion along their northern parts and a downward one along their southern parts.

Despite the fact that the quaternary fault tectonics is represented almost entirely by normal faults, it is debatable whether tensional forces caused it (as proposed, among others, by Leeder and Gawthorpe, 1987; Collier, 1990; Doutsos and Piper, 1990; Leeder *et al.*, 1991;) as the rotation effects observed in Isthmus of Corinthos suggest. Undoubtedly, the faults of the northern Peloponnessos, if examined separately, are due to gravity tectonics; the question lies, however, in identifying the primary genetic mechanism and in being able to recognize the tensional or compressive character of the deformation-generating forces.

Thus, if one wishes to explain the neotectonic-quaternary evolution of the Gulf, one should take into account the following:

- The uplift of the northern part of Peloponnessos and the region of Iti (southern part of Sperchios basin).
- The varying degree of uplift throughout the Giona – Iti dipole to such a degree, so that erosional surfaces of the same age, i.e. Calyvia surface, show a southern dip.

- ❑ The occurrence of Neogene beds in the northern Peloponnessos and their absence from the southern Sterea Hellas.
- ❑ The asymmetry observed in the morphology of the sea floor of the Gulf.
- ❑ The morphological asymmetry of the northern and southern coast.
- ❑ The presence of the normal antithetic faults in the northern part of Peloponnessos
- ❑ The mean WNW-ESE direction of the Gulf, which is determined by the *en échelon* arrangement of the E-W trending “marginal” faults.
- ❑ The vertical throw of the faults, as well as the thickness of sediments increases towards the East.

All these, combined with the segmented form of marginal fault zones and depocentres point to the conclusion that the central part of the Gulf is still at an older stage of evolution than the eastern one; the structure and evolution of the Gulf may not be the result of axial extension. The mechanics should involve more complex procedures, including those of a composite stress field (transtension and transpression) and the interaction between brittle and plastic deformation.




Fig. 6.1 Asymmetry of sea-floor morphology in Corinthiakos Gulf.




Fig. 6.2 Simplified tectonic map of western Corinthiakos Gulf showing bathymetry (continuous lines), extent and depth of continental shelf (dotted lines –depth in m.) and main active faults (hachured lines) (After Papanikolaou *et al.*, 1997).

Delphi

I. Mariolakis, I. Fountoulis & H. Kranis

*University of Athens, Division of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84 Athens*

Greece's numerous archaeological sites can provide excellent case studies for various subjects related to engineering geology, tectonics and morphotectonics, and historical seismicity, coupled with the intervention of human activity in the course of centuries.

A suitable case study is the ancient town of Delphi, located 250 km west of Athens in the southern slopes of Mt. Parnassos, and overlooking the northern margin of the Gulf of Corinth. It is one of the most important archaeological sites in Greece, where the most famous oracle of the ancient world was established.

In the geological map of **Fig. 7.1** two types of formations, the post-alpine and the alpine are shown.

The post-alpine formations are mainly represented by scree and talus cones, which cover the major part of the archaeological site. There is a very close relationship between these formations and (i) the very intense morphological slopes and (ii) the successive reactivations of the Arachova – Delfi fault zone in the northern border of the archaeological site (**Figs. 7.1, 7.2**). Four generations of scree can be distinguished (**Fig. 7.2**). The first one is mainly represented by compact breccia and conglomerates with calcitic matrix. The second one is represented by breccia and conglomerates cemented by a pelitic material. The third generation consists of polymictic, unconsolidated material, including big blocks or fragments of limestones. The fourth generation is the result of the relative recent rockfalls and is represented by big blocks of limestones (from 0.5 to 10 m³).

The thickness of the post-alpine formation differs from place to place, depending on the paleo-morphology of the basement. So, in the northern part of the archaeological site (**Fig. 7.1**), it seems to be small (up to 2 m.), as it is proved by the very frequent basement outcrops. On the contrary, at the southern part it seems to be greater but it does not exceed 15 m.

The alpine formations belong to the Parnassos Unit and they are represented (**Fig. 7.1**) by the Paleocene flysch (alternations of sandstones, marls and pelites) and neritic Jurassic-Cretaceous limestones.

The alpine tectonics is characterized by a large-scale overturned isoclinal fold (**Fig.7.3**) resulting in the younger flysch underlying the older limestones, at the northern part of the archaeological site. The contact between the limestones and the flysch is mostly regular, while at places the flysch is thrust over the carbonates. A great number of thrusts and reverse faults can be also observed within the calcareous rockmass.

The neotectonics of the area is characterized by the existence of 1st order macro-blocks (horsts and grabens) separated by fault zones, as the 1st order Arachova – Delphi active one that juxtaposes the Parnassos horst in the north against the Itea graben in the south (**Fig. 7.4**).

Numerous second- or third-order faults cut the alpine and post-alpine formations. Some of these faults are active, and seem to be connected to the historical-to-present-day seismic activity of the area. As for the inactive faults, their surfaces are convex or concave, when they are small, and undulated when they are big, while some of them bear striations. On the other hand, the active faults of the area are almost planar and cut the inactive ones. They are usually accompanied by a zone of loose gouge.

The hydrogeological conditions of the area are defined and controlled by the alpine and post-alpine structure and contact or over-flow springs occur along the carbonate-flysch contact. The drainage of the area is accommodated by the Phaidriades and Erateini torrents, in the east and west, respectively. Note that the area is prone to destructive flash floods.

The seismic activity of the major area seems to have been very high since historical times. Ancient of Delphi was razed by the earthquake series of 600 BC. ($I_0=VIII-X$).

The probability of shallow, $M>6$ earthquake occurring until 2006 is $0.80 < p < 1.00$ and the expected intensity for a design period of 80 yrs. is VIII-IX (Papaioannou, 1986). The expected ground acceleration with a 63% probability is $250-275 \text{ cm/sec}^2$ for the next 50 years (Makropoulos, 1986).




Fig. 7.1 Geological sketch map of Delphi. 1: Scree; 2: Flysch; 3: Limestones; 4: Fault; 5: Rockfalls; 6: Rockfall-prone site; 7: Landslides; 8: Creep; 9: Subsidence; 10: Ancient remnants; 11: (1) Stadium; (2) Apollo Sanctuary; (3) Kastalia spring; (4) Gymnasium; (5) Tholos (after Mariolakos *et al.*, 1991).




Fig. 7.2 Four generations of scree in Delphi (after Mariolakos *et al.*, 1991).




Fig. 7.3 Schematic geological cross-section (after Mariolakos *et al.*, 1991).




Fig. 7.4 Local neotectonic setting (after Mariolakos *et al.*, 1991).

Minyan Ancient Flood Prevention Works (c. 1600 BC.)

I. Mariolakis, I. Fountoulis & H. Kranis

*University of Athens, Division of Dynamic Tectonic Applied Geology,
Panepistimioupolis Zografou, GR 157 84 Athens*

The flood prevention - drainage works in Lake Kopaida are the earliest in Europe, since their construction dates back to 1600 BC. The technique implemented by the ancient people of *Minyes* is the same as the one currently taught at the Higher Education Institutions all over the world. These works, together with the mines in *Lavrio*, are the most important archeological sites suitable for the investigation of the technology of the ancient times; however the mining facilities in *Lavrio* are much younger (c. 800 BC.).

The largest navigable canal of the ancient times (27 km long) was constructed and, according to Knauss, it was used for product transportation (**Figs. 8.1, 8.3**). The water level at some parts of it was 1.5 m. above the adjacent areas that were used for cultivation and the embankment was totally impermeable (**Fig. 8.2**). The concept of the construction of that floodway is totally different from the one applied by a British company at the beginning of this century; the latter bore significant drawbacks, and subsequently numerous problems arose.

After the construction of the canal by the *Minyes* the water level at some parts of it was 1.5 m. above the ground surface (**Fig. 8.2**), which was used for crops. The works were totally impermeable, and their stability was excellent. It should be noted that the present national road connecting Kastro and Orhomenos is partially constructed on the embankment.

These works are related to the stage(s) of climatic optimum(s) in the Holocene and any further archaeogeological study should be valuable for the estimation of climatic fluctuations during the last 10 ka. Events as Noah's flood, or the less familiar *Deukalion's* flood, reflect such climatic conditions, which are, in turn, related to flood-prevention works. Knowledge of such climatic changes is important for the contribution to the prediction of such future changes.

The disappearance of the *Minyes* and their civilization is connected to the destruction of their works by the thebian *Hercules*, who sealed the entrance to the sinkhole that funneled the water carried by their canal with a boulder. So the whole area of Kopaida was inundated again, destroying all flood-prevention works, crops, and finally all the towns built on the flanks of the valley. Obviously, the cause of the fall of the boulder was an earthquake. This boulder can nowadays be seen still blocking the entrance to the sinkhole and the site has been proposed for inclusion in the international *Geotope* list of UNESCO. We should also say that the whole area between the springs of Orhomenos to the town of Kastro (ancient *Kopes*) and the sinkhole of Neo Kokkino could be developed into a unique **Earliest European Flood Prevention Technology Park**.




Fig. 8.1 Kopaida: Hystero-Helladic period reconstruction of lake area (gray fill for winter high-stand, line fill for summer highstand). Also shown the flood prevention embankments (heavy lines) and location of ancient towns/hamlets (After Knauss, 1984)




Fig. 2 Cross-section of artificial embankment




Fig. 8.3 Schematic reconstruction of navigable canal of Melas river (after Knauss, 1984).

References

- Brooks, M., Clews, J.E., Melis, N.S., Underhill, J.R., 1988. Structural development of Neogene basins in western Greece. *Basin Research* **1**, 129-138.
- Collier, R.E., 1990. Eustatic and tectonic controls upon Quaternary coastal sedimentation in the Corinth Basin, Greece. *J. Geol. Soc., London*, vol. **147**, pp. 301-314.
- Dercourt, J., 1964. Contribution a l'etude geologique d'une secteur du Peloponnese septentrional. *Ann. Geol. D. Pays Hellen.*, **15**, 418p., Athenes cum lit..
- Doutsos, T., & Piper, D.J.W., 1990. Sedimentary and morphological evolution of the Quaternary, eastern Corinth rift, Greece: first stages of continental rifting. *Bull. Geol. Soc. Am.*, **102**, pp. 812-829.
- Drakopoulos, J., Delibasis, N., 1982. The focal mechanism of earthquakes in the major area of Greece for the period 1947-1981. *Univ. Athens, Seismological Laboratory, publ.*, No **2**, Athens.
- Freyberg, B. Y. 1973. Geologie des Isthmus von Korinth. Erlanger Geol. Abh. Heft 95, 183 seiten.
- Galanopoulos, a., 1972. Plate tectonics in the area of Greece as reflected in the deep focus seismicity. *Bull. Geol. Soc. Greece*, **IX/2**, p. 266-285.
- Hatzfeld, D., Pedotti, G., Hatzidimitriou, P., Makropoulos, K., 1990. The strain pattern in the western Hellenic Arc deduced from microearthquake survey. *Geophys. J.*, **101/1**, 181-202.
- Heezen, B., Ewing, M., Johnson, L.G., 1966. The Gulf of Corinth floor. *Deep Sea Research*, **13**, 381-411.
- Higgs, B., 1988. Synsedimentary structural controls on basin deformation in the Gulf of Corinth, Greece. *Basin Research*. **1**, 155-165.
- Kelletat, D., Schröder, B., 1975. Vertical displacement of Quaternary shorelines in the Peloponnesus, Greece. *Proc. Verb. CIESM Congr.*, Monaco.
- Laj, G., Jamet, M., Sorel, D., Valente, S.R., 1982. First paleomagnetic results from Miopliocene series of the hellenic sedimentary arc. *Tectonophysics*, **86**, 45-62.
- Le Pichon, X. & Angelier, J., 1979. The Hellenic Arc and Trench System: a key to the neotectonic evolution of the eastern Mediterranean area. *Tectonophysics*, v. 60, p. 142.
- Le Pichon, X. & Angelier, J., 1981. The Aegean sea. *Phil. R. Soc. London*, **A 300**, 357-372.
- Le Pichon, X. & Angelier, J., Sibuet, J.C., 1982. Plate boundaries and extensional tectonics. *Tectonophysics*, **81**, 239-256.
- Le Pichon, X., Lyberis, N., Angelier, J., Renard, V., 1982. Strain distribution over the east Mediterranean ridge: a synthesis incorporating new seabeam data. *Tectonophysics*, **86**, 243-274.
- Leeder, M.R., & Gawthorpe, R.L., 1987. Sedimentary models for extensional tilt block/half graben basins. In: Leeder, M.R., Dewey, J.F., & Hancock, P.L. eds *Cont. Ext. Tectonics. Geol. Soc., London, Sp. Publ.*, **28**, 139-152.
- Leeder, M.R., Seger, M.J., Stark, C.P., 1991. Sedimentation and tectonic geomorphology adjacent to major and inactive normal faults, southern Greece. *J. Geol. Soc., London*, **148**, 331-343.
- Makris, J. 1978. A geophysical study of Greece based on deep seismic soundings, gravity and magnetics. In Cloos, H., Roeder, R., Schmidt, K., (eds.), *Alps, Apennines, Hellenides*. Stuttgart E. Schweizerbartsche Verlagsbuchhandlung, 392-401.
- Makropoulos, K., G. Stavrakakis, S. Latoussakis, I. Drakopoulos, 1986. A comparative seismic hazard study for the area of Greece, *Proc. 2nd Int. Seminar on Earth. Progn.*, Berlin.
- Mariolakos, I. & Stiros, S., 1987. Quaternary deformation of the Isthmus and Gulf of Corinthos

- Greece. *Geology*, **15**, 225-228.
- Mariolakos, I. & Papanikolaou, D., 1981. The neogene basins of the Aegean Arc from the Paleogeographic and the Geodynamic point of view. *Proc. Int. Symp. Hell. Arc and Trench HEAT*, t. **I**, p. 383-399, Athens.
- Mariolakos, I. & Papanikolaou, D., 1987. Deformation pattern and relation between deformation and seismicity in the Hellenic arc. *Bull. Geol. Soc. Greece*, **XIX**, p. 59-76 (in Greek).
- Mariolakos, I., 1975. Thoughts and viewpoints on certain problems of the Geology and tectonics of Peloponnesus Greece. *Ann. Geol. Pays Hellen.*, **27**, p. 215-313 (in Greek).
- Mariolakos, I., Fountoulis, I., Logos, E., Lozios, S., 1989. Surface faulting caused by the Kalamata Greece earthquake 1391986. *Tectonophysics*, **163**, 197-203.
- Mariolakos, I., Fountoulis, I., Logos, E., Lozios, S., 1991. Methods to study torsional neotectonic deformation: the case of Kalamata area SW Peloponnesus, Greece. In Chen Qingxuan (Ed.) *Proc. Regional Crustal Stability and Geological Hazards, IGCP project 250*, vol. **3**, 15-21, *Inst. Geomechanics, Chinese Academy of Geological Sciences CAGS*, Beijing.
- Mariolakos, I., Fountoulis, I., MarcopoulouDiacantoni, A., Mirkou, M.R., 1994. Some remarks on the kinematic evolution of Messinia Province SW Peloponnesus, Greece during the Pleistocene based on Neotectonic, Stratigraphic and Paleoecological observations. *Munster. Forsch. Geol. Palaont.*, **76**, 371-380, Munster.
- Mariolakos, I., Logos, E., Lozios, S., Nassopoulou, S., 1991. Technicogeological observations in the Ancient Delphi area Greece. In: AlmeidaTeixeira, R. Fantechi, R. Oliveira, A. Gomes Coelho (Eds.), *Prevention and control of landslides and other mass movements, Report EUR 12918 EN*, 273-283.
- Mariolakos, I., Papanikolaou, D., Lagios, E., 1985. A neotectonic geodynamic model of Peloponnesus based on; morphotectonics, repeated gravity measurements and seismicity, *Geol., Jb.*, Band **50**, p. 317.
- Maul, Ot., 1921. Beitrage zur Morphologie des Peloponnes und des sudlichen Mittelgriechenlands. *Geogr. Abhandl.*, **10**, fasc. 3, 120 p. Leipzig.
- Mc Kenzie, D.P., 1970. Plate tectonics in the Mediterranean region. *Nature*, **226**, 239-243.
- Mc Kenzie, D.P., 1972. Active tectonics of the Mediterranean region. *Geoph. J. R. Astron. Soc.*, v. **30**, 109-185.
- Mc Kenzie, D.P., 1978. Active tectonics of the AlpineHimalyan Belt: the Aegean Sea and surrounding regions. *Geophys. J. R. Astron. Soc.*, **55**, 217-254.
- Mercier, J.L., 1979. Signification neotectonique de l' Arc Egeen. Une revue des idees. *Rev. Geol. Dyn. Geogr. Phys.*, **1**, 5-15.
- Mercier, J.L., Delibasis, N., Gautier A., Jarrige, J.J., Lameille, F., Philip, H., Sbrier, M., Sorel, D., 1979. La Neotectonique de l' Arc Egeen: *Revue de Geographie Physique et de Geologique Dynamique*, v. **21**, fasc 1, p. 67-92.
- Papaioannou, C., 1986. Seismic hazard assessment and longterm earthquake prediction in southern Balkan region, *Proc. 2nd Int. Seminar on Earth. Progn.*, Berlin.
- Papanikolaou, D., 1986. Geology of Greece. *Eptalofos publ.* 240 p., Athens.
- Papanikolaou, D., Chronis, G., Lykousis, V., Sakellariou, D., Papoulia, I., 1997. Submarine Neotic Structure of W. Korinthiakos Gulf and Geodynamic phenomena of the Egion Earthquake. *Proceedings, 5th Panhellenic Symposium of Oceanography and Fishery*, **1**, 415-418 (in Greek).
- Papanikolaou, D., Lykousis, V., Chronis, G., Pavlakis, P., 1988. Acomparative study of neotectonic basins across the Hellenic arc: the Messiniakos, Argolikos, Saronikos and Southern Evoikos Gulfs, *Basin Research*, **1**, 167-176.
- Paquin, C., Bloyet, J., Angelidis, C., 1984. Tectonic stresses on the boundary of the Aegean domain: insitu measurements by overcoring. *Tectonophysics*, **110**, 145-150.

- Paquin, C., Froidevaux, C., Bloyet, J., Ricard, Y., Angelidis, C., 1982. Tectonic stresses on the mainland of Greece: insitu measurements by overcoring. *Tectonophysics*, 86, 17-26.
- Philippson, A., 1892. *Der Peloponnes*. Verlag Friedlander, 642s, Berlin.
- Philippson, A., 1930. *Beitrage zur morphologie Griechenlands*. Geogr. Abhndl., 3, Dritte Reih., 93s., Stuttgart.
- Ritsema, A. R., 1974. The earthquake mechanism of the Balkan region. *Roy. Netherlands Meteorol. Inst. Sci., Rep. 744*, p. 136.
- Sebrier, M., 1977. *Tectonique Recente d' une transversale a l' Arc Egeen* Ph.D. thesis, Paris, Universite de Paris XI, 76 p..
- Sweeting, M.M. 1967. Report on the geomorphogeny of the ParnassosGhiona region of Greece. *I.G.S.R. Geol. Rec. 41a*, 25 p., Athens.
- Tselentis, G.A., Makropoulos, K., 1986. Rates of crustal deformation in the Gulf of Corinth central Greece as determined from seismicity. *Tectonophysics*, 124, 55-66.